

Optics, Acoustics, and Stress in a Nearshore Bottom Nepheloid Layer (OASIS)

John H. Trowbridge
Woods Hole Oceanographic Institution
Woods Hole, MA 02543
phone: 508-289-2296 fax: 508-457-2194 e-mail: jtrowbridge@whoi.edu

Emmanuel Boss
School of Marine Sciences
5741 Libby Hall
University Of Maine
Orono, ME 04469-5741
phone: (207) 581-4378 fax: (207) 581-4388 email: emmanuel.boss@maine.edu

Paul S. Hill
Department of Oceanography
Dalhousie University
Halifax, Nova Scotia, CANADA B3H 4J1
phone: (902) 494-2266 fax: (902) 494-3877 email: paul.hill@dal.ca

Timothy G. Milligan
Fisheries and Oceans Canada
Bedford Institute of Oceanography
1 Challenger Drive
Dartmouth, Nova Scotia, CANADA B2Y 4A2
phone: (902) 426-3273 fax: (902) 426-6695 email: milligant@mar.dfo-mpo.gc.ca

Award number: N000140410221
<http://www.whoi.edu/mvco/projects/projects.html>

LONG-TERM GOALS

To quantify and understand the effects of aggregation dynamics on the distribution of particles in the bottom boundary layer, and to understand how the properties of particles (composition, shape, and internal structure) affect their optical and acoustical properties.

OBJECTIVES

- Obtain a set of acoustical and optical measurements that determines the evolution of the particle size distribution.
- Obtain concurrent velocity, temperature and conductivity measurements sufficient to determine the fluid dynamical environment within which the particle size distribution evolves.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2009		2. REPORT TYPE		3. DATES COVERED 00-00-2009 to 00-00-2009	
4. TITLE AND SUBTITLE Optics, Acoustics, and Stress in a Nearshore Bottom Nepheloid Layer (OASIS)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Woods Hole Oceanographic Institution, Woods Hole, MA, 02543				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

- Evaluate and improve a state-of-the-art model of particle dynamics by comparing model calculations with measurements.

APPROACH

The approach is to obtain measurements that will permit comparisons of the optical and acoustical signatures of suspended particles and inferences of the particle size distribution and its temporal evolution, concurrently with fluid dynamical measurements that determine the flow field within which the particles evolve. The instrumentation is mounted on bottom tripods and includes a 9-wavelength optical attenuation and absorption meter (WetLabs ac-9, with automated regular dissolved measurement for calibration independent particulate measurements), a LISST-100 laser diffraction particle sizer (Agrawal & Pottsmith 2000), a digital floc camera (DFC) (Curran et al. 2002b), a Tracor Acoustic Profiling System (TAPS, Holliday 1987), and an array of SonTek/YSI acoustic Doppler velocimeters (ADV) . Near-simultaneous measurements with and without a filter assure high-quality particulate spectral absorption and attenuation measurements with the ac-9. The LISST-100 and floc camera together provide particulate size distributions from 2.5 micrometers to 1 centimeter. The TAPS obtains range-gated, vertical profiles of acoustical backscatter intensity at a range of frequencies between 0.3 and 3.0 MHz. The TAPS and ADVs produce acoustical measurements over a wide range of frequencies that can be used to generate particle size distributions (Holliday, 1987; Hay and Sheng, 1992). The combined optical and acoustical measurements will provide a comprehensive description of the suspended particles near the seabed. The velocity measurements obtained from the ADVs will provide direct-covariance estimates of Reynolds stress and inertial-range estimates of the dissipation rate for turbulent kinetic energy (Trowbridge 1998, Trowbridge & Elgar 2001, Shaw & Trowbridge 2001, Trowbridge & Elgar 2003).

The analysis includes estimation of Reynolds stress, dissipation rate, and eddy diffusivity; estimation of particle size distribution and concentration from the DFC and LISST-100; and estimation the optical and acoustical properties of the water column from analysis of the ac-9, TAPS, and ADV data. The analysis focuses on evaluation and improvement of a one-dimensional (vertical), time-dependent model of the particle size and concentration fields and the accompanying optical and acoustical properties. The model includes the effects of sediment resuspension by bottom shear stresses produced by waves and currents, vertical transport of suspended particles by turbulence, gravitational settling of particles, and particle aggregation and disaggregation.

WORK COMPLETED

Three sets of measurements have been obtained, the first during August-September 2004, the second during August-September 2005, and the third during August-September 2007. A fourth set of measurements is currently underway (September 2009). All four sets of measurements have occurred at the Martha's Vineyard Coastal Observatory (MVCO), which is operated by the Woods Hole Oceanographic Institution (WHOI). The MVCO, off the southern coast of Martha's Vineyard, Massachusetts, is a cabled observatory consisting of a shore station, a meteorological mast on the beach, a seafloor node at a water depth of 12 m, and an air-sea interaction tower (ASIT) at a water depth of 15 m. Atmospheric measurements are obtained routinely at the meteorological mast and the ASIT. Routine oceanic measurements of temperature, salinity and velocity are obtained at the 12-m node and the ASIT. The 2004 measurements for this study were obtained near the ASIT and the 2005

and 2007 measurements were obtained near the 12-m node. The 2007 program included measurements obtained from a profiler deployed by WetLabs.

RESULTS

Analysis of OASIS measurements has focused on characterization of the near-bottom flow environment in which the suspended particles exist and evolve. The flow characterization includes quantitative estimates of the bottom stress, dissipation rate, and mean shear. The measurements indicate a production-dissipation balance in the turbulent kinetic energy equation and excellent agreement with the Prandtl-Karman law of the wall, as expected in an un-stratified shear flow (Figure 1). A primary result of the analysis is estimates of the stress associated with the hour-averaged current and the standard deviation of the stress produced by surface gravity waves (Figure 2). The analysis indicates good agreement between measured hour-averaged stresses and corresponding stresses estimated from a boundary layer model, and much larger oscillatory stresses produced by surface gravity waves. Work in progress addresses the relationship between the stresses and the characteristics and concentration of the suspended particles. Additional work has addressed the interaction between turbulence and suspended organisms (Jumars et al. 2009) and the effect of breaking waves and Langmuir circulations on water-column turbulence (Gerbi et al. 2009, Kukulka et al. 2009).




Figure 1. A test of the relationship $(\kappa z \varepsilon)^{2/3} = |\langle uw \rangle|$ based on the 2007 OASIS measurements at the Martha's Vineyard Coastal Observatory. Here $\kappa \approx 0.40$ is the Karman constant, z is the distance above the bottom, ε is the hour-averaged energy dissipation rate per unit mass, and $-\rho \langle uw \rangle$ is the turbulent Reynolds shear stress, where ρ is the fluid density, u and w are the streamwise and vertical turbulent velocity fluctuations, and brackets denote an hour average. The relationship $(\kappa z \varepsilon)^{2/3} = |\langle uw \rangle|$ holds if the hour-averaged velocity varies logarithmically with height and if shear production of turbulent kinetic energy balances dissipation, as in a classical unstratified wall layer.




Figure 2. *Stresses estimated from the 2007 measurements at the Martha’s Vineyard Coastal Observatory. The directly measured hour-averaged Reynolds shear stress $-\rho \langle uw \rangle$ agrees well with the drag-law estimate $c_d \rho |U| U$ inferred from the measured burst-averaged velocity U by means of an empirically determined drag coefficient c_d . The standard deviation ρu_{wm}^2 of the oscillatory bottom shear stress produced by surface waves is inferred from a wave-current interaction model and is much larger than the hour-averaged stress $-\rho \langle uw \rangle$, which is associated with wind-driven and tidal currents. The wave-induced stress is primarily responsible for suspending particles and controlling the size of water-borne flocs.*

IMPACT/APPLICATIONS

Operational seagoing systems often depend on optical and acoustical properties of suspended particles in the water column. Understanding the processes that regulate the particle characteristics and understanding the optical and acoustical signatures of suspended particles are essential in order to predict the performance of these operational systems.

RELATED PROJECTS

Mechanisms of Fluid-Mud Interactions under Waves, an ONR-funded MURI project aimed at understanding the dissipation of surface gravity waves over muddy seafloors. This combined field, laboratory, numerical, and theoretical study has been undertaken by S. J. Bentley (Memorial

University), R. A. Dalrymple (Johns Hopkins University), G. C. Kineke (Boston College), C. C. Mei (Massachusetts Institute of Technology), P. Traykovski (Woods Hole Oceanographic Institution), J. H. Trowbridge (WHOI), and D. Yue (MIT). Companion field and analysis studies are being carried out by S. Elgar (WHOI), B. Raubenheimer (WHOI), T. Herbers (Naval Postgraduate School), and A. Sheremet (University of Florida). A focus of the WHOI, Boston College, and Memorial University field work is quantitative imaging and interpretation of the near-bottom mud dynamics that control energy dissipation.

REFERENCES

1. Agrawal, Y. C. and H. C. Pottsmith, 2000: Instruments for particle size and settling velocity observations in sediment transport. *Marine Geology*, 168: 89-114.
2. Curran, K. J., P. S. Hill, and T. G. Milligan. 2002. The role of particle aggregation on size-dependent deposition of drill mud. *Continental Shelf Research*, 22: 405-416.
3. Holliday, D. V. 1987. Acoustic Determination of Suspended Particle Size Spectra. Proceedings of a Specialty Conference on Acoustical Measurements.
4. Shaw, W.J. and J.H. Trowbridge, 2001. The direct estimation of near-bottom turbulent fluxes in the presence of energetic wave motions. *Journal of Atmospheric and Oceanic Technology*, 18: 1540-1557.
5. Trowbridge, J.H. and S. Elgar, 2001. Turbulence measurements in the surf zone. *Journal of Physical Oceanography* 31: 2403-2417.
6. Trowbridge, J.H. and S. Elgar, 2003. Spatial scales of stress-carrying nearshore turbulence. *Journal of Physical Oceanography*, 33:1122-1128.

PUBLICATIONS

1. Gerbi, G. P., J. H. Trowbridge, E. A. Terray, A. J. Plueddemann, and T. Kukulka, 2009. Observations of turbulence in the ocean surface boundary layer: energetics and diffusivity. *Journal of Physical Oceanography*, 39, 1077-1096.
2. Jumars, P. A., J. H. Trowbridge, E. Boss, and L. Karp-Boss, 2009. Turbulence-plankton interactions: a new cartoon. *Marine Ecology*, 30: 133-150.
3. Kukulka, T., A. J. Plueddemann, J. H. Trowbridge, and P. P. Sullivan, 2009. Significance of Langmuir circulation in upper ocean mixing: comparison of observations and simulations. *Geophysical Research Letters*, 36, L10603, doi:10.1029/2009GL037620.